

CONTENT

Content Page	A
Editorial <i>J. Hinchliffe</i>	B
Age estimation in an Indian population using pulp/tooth volume ratio of mandibular canines obtained from cone beam computed tomography. <i>N. Jagannathan, P. Neelakantan, C. Thiruvengadam, P. Ramani, P. Premkumar, A. Natesan, J.S. Herald, H. U. Luder.</i>	1 - 6
Stature estimation by Carrea's Index and its reliability in different types of dental alignment. <i>L. Lima, Y. da Costa, R. Tinoco, P. Rabello, E. Daruge Junior.</i>	7 - 13
Rugae patterns as an adjunct to sex differentiation in forensic identification. <i>A. Saraf, S. Bedia, A. Indurkar, S. Degwekar, R. Bhowate.</i>	14 - 19
Establishing the reliability of palatal rugae pattern in individual identification (following orthodontic treatment). <i>D. Shukla, A. Chowdhry, D. Bablani, P. Jain, R. Thapar.</i>	20 - 29
Inter canine distance in the analysis of bite marks: A comparison of human and domestic dog dental arches. <i>S.V. Tedeschi-Oliveira, M. Trigueiro, R.N. Oliveira, R.F.H. Melani.</i>	30 - 36
Sexual dimorphism in the permanent maxillary first molar: a study of the Haryana population (India). <i>V. Sonika, K. Harshaminder, G.S. Madhushankari, J.A. A Sri Kennath.</i>	37 - 43
Case reports and background: Difficulties with identification – Sweden. <i>I. Dawidson</i>	44 - 50
Book Revue <i>J. A. Hinchliffe</i>	51